

1 **Toll-like Receptor 2 Costimulation Potentiates the Antitumor**
2 **Efficacy of CAR T Cells**

3 **Authors:**

4 Yunxin Lai^{1,2}†, Jianyu Weng³†, Xinru Wei^{1,2}, Le Qin^{1,2}, Peilong Lai³, Rucong Zhao^{1,}
5 ², Zhiwu Jiang^{1,2}, Baiheng Li^{1,2}, Simiao Lin^{1,2}, Suna Wang^{1,2}, Qiting Wu^{1,2}, Zhaoyang
6 Tang^{4,5}, Pentao Liu⁶, Duanqing Pei^{1,2}, Yao Yao^{1,2}, Xin Du^{3*}, Peng Li^{1,2,7*}

7 **Affiliations:**

8 ¹Key Laboratory of Regenerative Biology, South China Institute for Stem Cell Biology
9 and Regenerative Medicine, Guangzhou Institutes of Biomedicine and Health, Chinese
10 Academy of Sciences, Guangzhou, 510530, China;

11 ²Guangdong Provincial Key Laboratory of Stem Cell and Regenerative Medicine,
12 South China Institute for Stem Cell Biology and Regenerative Medicine, Guangzhou
13 Institutes of Biomedicine and Health, Chinese Academy of Sciences, Guangzhou,
14 510530, China;

15 ³Department of Hematology, Guangdong General Hospital/Guangdong Academy of
16 Medical Sciences, Guangzhou 510080, Guangdong, China;

17 ⁴Guangdong Zhaotai InVivo Biomedicine Co. Ltd., Guangzhou, 510000, China;

18 ⁵Hunan Zhaotai Yongren Medical Innovation Co. Ltd., Changsha, 410000, China;

19 ⁶Wellcome Trust Sanger Institute, Hinxton, Cambridge CB10 1HH, England, UK;

1 ⁷Department of Abdominal Surgery, Affiliated Cancer Hospital & Institute of
2 Guangzhou Medical University of Guangzhou Medical University, Guangzhou
3 Medical University, Guangzhou, Guangdong, 510095, China.

4 *Correspondence to: Peng Li, PhD, email: li_peng@gibh.ac.cn; Xin Du, MD/PhD,
5 email: miyadu@hotmail.com.

6 **Footnotes:** † indicates co-first authorship.

7 **One Sentence Summary:**

8 The innate immune receptor TLR2, incorporated into CARs, enhanced the efficacy of
9 CAR T cells in eliminating leukemia and solid tumors.

10

1 **Abstract**

2 Chimeric antigen receptor (CAR) T cell immunotherapies have shown unprecedented
3 success in treating leukemia but limited clinical efficacy in solid tumors. Here, we
4 generated 1928zT2 and m28zT2, targeting CD19 and mesothelin, respectively, by
5 introducing the Toll/interleukin-1 receptor (TIR) domain of Toll-like receptor 2 (TLR2)
6 to 1928z and m28z. T cells expressing 1928zT2 or m28zT2 showed improved
7 expansion, persistency and effector function against CD19⁺ leukemia or mesothelin⁺
8 solid tumors respectively in vitro and in vivo. In a patient with relapsed B cell acute
9 lymphoblastic leukemia, a single dose of 5×10^4 /kg 1928zT2 T cells resulted in robust
10 expansion and leukemia eradication and led to complete remission. Hence, our results
11 demonstrate that TLR2 signaling can contribute to the efficacy of CAR T cells. Further
12 clinical trials are warranted to establish the safety and efficacy of this approach.

13

14

15

1 Introduction

2 The success of the adoptive transfer of CD19-specific chimeric antigen receptor
3 (CAR) T cells represents a major step in cancer immunotherapy. Both CD3 ζ and
4 costimulatory domains ¹ are needed to generate effective CARs that provide the
5 signals necessary for full activation, expansion and survival of CAR T cells. The
6 incorporation of costimulatory molecules CD28 ², 4-1BB ^{3,4}, OX-40 ⁵, CD27 ⁶ and
7 ICOS ⁷ significantly improve the expansion, cytokine production, in vivo persistence
8 and antitumor efficacy of CAR T cells. Moreover, the efficacy of CAR T cells for
9 solid tumors is substantially poorer than that for leukemia ^{8, 9}. Given the few
10 costimulatory molecules that have been used in the design of CARs, novel
11 intracellular signals are needed to improve their efficacy so as to effectively treat not
12 only leukemia but also solid tumors.

13 The intracellular signaling domains of CARs mainly derive from the
14 costimulatory receptors expressed on T cells that are critical for their adaptive
15 immune responses. Nevertheless, the innate immune receptor Toll-like receptor 2
16 (TLR2) is highly expressed on activated and memory T cells, and the TLR2 signaling
17 increases T cell expansion and cytokine production ¹⁰⁻¹³, lowers the activation
18 threshold for costimulatory signals delivered by antigen presenting cells (APCs),
19 facilitates the generation of memory CD8 T cells ¹⁴⁻¹⁶, and directly triggers Th1
20 effector functions ¹⁷. In addition, TLR2-mediated signaling can abolish the
21 suppressive capacities of regulatory T (Treg) cells¹⁸. Furthermore, T cell
22 hyporesponsiveness with impaired IFN- γ and IL-2 production can partially be

accounted for by diminished expression of TLR2 on T cells¹⁹. Strikingly, TLR2 ligands induce the regression of established leukemia and solid tumors through compromising the suppressive function of Foxp3⁺ Treg cells and enhancing the cytotoxicity of tumor-specific T cells^{20, 21}. TLR2 can form heterodimers with TLR1 or TLR6 to initiate a proinflammatory response^{22, 23}. However, TLR2 may also form weak homodimers, which allow it to directly bind to MyD88²⁴. Therefore, we proposed that TLR2 signaling could potentiate the antitumor activity of CAR T cells.

To test our hypothesis, we generated CARs targeting CD19 (1928z) and mesothelin (m28z) by linking scFv with CD28 and CD3 ζ signaling domains and then generated 1928zT2 or m28zT2 by adding the TIR domain of TLR2 to the 3' end of CD3 ζ . We compared the functions of CAR T cells with or without the TLR2 signaling domain and found that the addition of a TLR2 domain led to enhanced antitumor responses for both leukemia and solid tumors. To test the efficacy and safety of TLR2 construct in humans, we conducted a pilot trial in a patient with relapsed B cell acute lymphoblastic leukemia (B-ALL) with 1928zT2 T cells and found that a single dose (5 \times 10⁴/kg) of 1928zT2 T cells led to complete remission. Our data revealed that the innate immune receptor TLR2, in addition to canonical costimulatory receptors, can improve the antitumor efficacy of CARs for both leukemia and solid tumors.

Results

TLR2 enhanced the effector functions of CD19-specific CAR T cells in vitro

The CD19-specific second-generation CARs have generated exciting results in

1 clinical trials ²⁵⁻³⁰. We used 1928z, a CAR-linking FMC63-scFv, CD28
 2 transmembrane and endodomain, and the CD3 ζ signaling domain as benchmarks to
 3 evaluate TLR2 function in CARs. The TIR domain (amino acid 639-784) of TLR2
 4 was appended to the CD3 ζ signaling domain. Lentiviral vectors that contained GFP
 5 (negative control), 1928z and 1928zT2 (Supplementary Figure 1a) were transfected
 6 into T cells to generate GFP T, 1928z T and 1928zT2 T cells, respectively. The
 7 transduction efficiencies were measured by the percentages of GFP⁺ cells
 8 (Supplementary Figure 1b). To confirm the role of GFP in representing CAR
 9 expression, we detected the coexpression of GFP and anti-CD19 CAR
 10 (Supplementary Figure 1c). K562, K562-CD19, NALM6 and REH cells were tagged
 11 with both GFP and luciferase (GL). K562-CD19-GL, NALM6-GL, and REH-GL cells,
 12 but not K562-GL cells, were specifically killed by 1928z T and 1928zT2 T cells, and
 13 notably, the killing percentages of 1928zT2 T cells were higher than those of 1928z T
 14 cells at low effector to target (E/T) ratios (Figure 1a). Expression of *GZMB*, *PRF1* and
 15 *IFNG* were higher in 1928zT2 T cells compared to 1928z T cells upon stimulation by
 16 NALM6 cells (Supplementary Figure 2). Consistently, 1928zT2 T cells secreted
 17 significantly higher amounts of IL-2, IFN- γ and GM-CSF, but a lower level of TNF- α
 18 (Figure 1b). Interestingly, suppression of TNF- α secretion from macrophages by
 19 TLR2 agonist has been reported ³¹. In addition, the capacities of 1928zT2 T cells to
 20 expand upon serial stimulations and to continuously eliminate target cells were higher
 21 than 1928z T cells (Figure 1c). Moreover, the killing capacity of 1928zT2 T cells was
 22 enhanced compared with 1928BBz T cells (Supplementary Figure 3). Taken together,

1 the incorporation of the TLR2 signaling domain into 1928z led to enhanced
2 anti-tumor effector functions in vitro.

3 *1928zT2 T cells exhibited enhanced antitumor efficacy in vivo*

4 Next, the killing capacities of 1928z T and 1928zT2 T cells were compared in
5 cell line-derived xenografts. Immunodeficient NSI mice ³² were intravenously
6 injected with 1×10^6 NALM6-GL cells followed by two infusions of 5×10^6 GFP T,
7 1928z T or 1928zT2 T cells. Bioluminescence imaging (BLI) showed that a reduced
8 tumor burden was present in mice infused with 1928zT2 T cells compared with 1928z
9 T cells (Figure 2a). Consistently, the percentages of NALM6-GL cells in the
10 peripheral blood mononuclear cells (PBMC) were significantly lower in mice treated
11 with 1928zT2 T cells compared with 1928z T cells (Figure 2b). In addition, the mice
12 treated with 1928zT2 T cells exhibited delayed weight loss (Supplementary Figure
13 S4a). Furthermore, mice treated with 1928zT2 T cells showed enhanced secretion of
14 IL-2, IFN- γ and GM-CSF, but not TNF- α , in serum (Supplementary Figure 4b). To
15 confirm the improved efficacy of the TLR2 construct in terms of survival, 2×10^5
16 NALM6 cells, and two days later, 2×10^6 GFP T, 1928z T, 1928zT2 T or no T cells,
17 were intravenously injected into each NSI mice (n=10). 1928zT2 T cells improved the
18 survival of NALM6 burdened mice better than 1928z T cells (Figure 2c). The
19 percentages of NALM6 cells and T cells in BM when mice died (Supplementary
20 Figure 4c) indicated that the major cause of death was high NALM6 burden for mice
21 treated with no T, GFP T and 1928z T cells, but was graft-versus-host disease (GVHD)
22 for mice treated with 1928zT2 T cells as severe symptoms such as loss of hair and

weight were observed. Thus, 1928zT2 T cells were more potent than 1928z T cells at killing NALM6-GL cells in vivo.

Furthermore, the efficacies of 1928z T and 1928zT2 T cells were compared in patient-derived xenograft (PDX) models. B-ALL cells were eliminated by 1928zT2 T cells, but not by 1928z T cells, and the percentages of total human T cells and CAR T cells were higher in 1928zT2 group compared with 1928z group (Supplementary Figure 5a). Due to the less aggressiveness of primary B-ALL cells, all the B-ALL burdened mice including the GFP T group survived until when the mice were culled for analysis on day 78, and splenomegaly was found only in the GFP and 1928z groups, but not in the 1928zT2 group (Supplementary Figure 5b). Collectively, these results demonstrated that 1928zT2 T cells had significantly higher anti-leukemia efficacy than 1928z T cells in vivo.

TLR2 costimulation upregulated genes associated with cell adhesion, synaptic transmission, T cell migration and mTOR signaling

To decipher the molecular mechanisms of TLR2 costimulation, we performed RNA sequencing to compare the transcriptomes of 1928z T and 1928zT2 T cells with or without stimulation by NALM6 cells. Approximately 11013 of the 33382 RefSeq genes were detectably expressed in all four populations. We identified the genes specifically upregulated in stimulated 1928zT2 T cells and found that most of these genes were related to cell adhesion and synaptic transmission (Figure 3a). Quantitative real-time PCR confirmed the upregulation of genes associated with cell adhesion and synaptic transmission, along with two genes *IRF3* and *IRF7* downstream

1 of the TLRs pathway (Figure 3b). TLR2 signaling has been reported to enhance
2 mTOR signaling and expression of T-bet ³³. Here, we also detected elevated
3 transcription of *T-bet*, *mTOR*, *4E-BP1* and *P70S6K* (Figure 3b). Matrix
4 metalloproteinases (MMPs) have been shown to play critical roles in T cell activation
5 ³⁴ and migration into tissues ³⁵. TLR2 incorporation also induced the upregulation of
6 MMP2 and MMP24 expression in 1928zT2 T cells (Figure 3a and 3b). Therefore, the
7 migration capacities of GFP T, 1928z T and 1928zT2 T cells were compared and
8 results indicated that 1928zT2 T cells transmigrated the matrigel more efficiently than
9 1928z T cells (Figure 3c). These results suggest that TLR2 incorporation enhances the
10 antitumor efficacy of CAR T cells partly through improved adhesion, synaptic
11 transmission and migration capacities, and enhanced T-bet expression through mTOR
12 pathway.

13 *TLR2 costimulation enhanced antitumor efficacy of mesothelin-specific CAR T cells*

14 Mesothelin is highly expressed in various solid tumors including lung
15 adenocarcinoma cells ³⁶, and clinical trials of mesothelin-targeting CAR T cells are
16 ongoing (such as NCT02414269 and NCT02930993). Thus, to assess the effect of
17 TLR2 costimulation on CAR T cells targeting solid tumors, we incorporated TLR2
18 into a CAR specific to mesothelin and generated GFP T, m28z T and m28zT2 T cells
19 through lentiviral transfection (Supplementary Figure 6a and 6b). m28zT2 T cells
20 showed enhanced killing capacity against mesothelin⁺ A549-GL cells compared with
21 m28z T cells (Figure 4a). m28zT2 T cells secreted more IL-2, IFN- γ and GM-CSF,
22 but reduced TNF- α compared with m28z T cells upon stimulation by A549 cells

1 (Figure 4b). The expansion of m28zT2 T cells during repetitive stimulation was
2 improved compared with m28z T cells, and the potency of m28zT2 T cells at killing
3 A549 cells were significantly higher than m28z T cells (Supplementary Figure 7a).
4 Meanwhile the exhaustion markers on T cells after repetitive coculture were
5 compared, and the expression of TIM-3, but not PD-1 or LAG-3, was lower on
6 m28zT2 T cells relative to m28z T cells (Supplementary Figure 7b). These results
7 indicated that TLR2 incorporation in mesothelin-specific CAR also improved its
8 effector functions in vitro.

9 We further compared the antitumor efficacy of m28zT2 T, m28z T and GFP T
10 cells in subcutaneous A549-GL xenografted mice. Tumors from m28zT2 group were
11 smaller than those from the blank, GFP and m28z groups (Figure 4c). Moreover, mice
12 treated with m28zT2 T cells produced more cytokines IL-2, IFN- γ and GM-CSF, but
13 not TNF- α , than those treated with m28z T cells (Supplemental Figure 8). Since
14 metastasis is a major hurdle in the treatment of solid tumors, we compared the
15 efficacies of m28z T and m28zT2 T cells against metastasized tumors through the
16 blood in intravenous A549-GL xenografted mice. The BLI results showed that
17 m28zT2 T cells efficiently inhibited the growth of A549-GL cells in NSI mice;
18 however, surprisingly, the mice grew a greater burden of tumors when treated with
19 m28z T cells than the mice that were injected with GFP T cells (Supplementary Figure
20 9a). The histologic analysis confirmed that the tumor cells in the uterus of the mice
21 from the m28z group were HLA⁺ A549-GL cells (Supplementary Figure 9b). These
22 results indicated that TLR2 incorporation could enhance the efficacy of

1 mesothelin-specific CAR T cells against solid tumors.

2 *Clinical trial of 1928zT2 T cells in a patient with relapsed and refractory B-ALL*

3 We conducted a pilot clinical trial in a patient with relapsed and refractory
4 B-ALL (NCT02822326). 1928zT2 T cells were produced and well characterized
5 phenotypically and functionally (Supplementary Figure 10a-f). The patient received
6 two days of treatment with Fludarabine (30 mg/m²/day) and cyclophosphamide (300
7 mg/m²/day) for lymphocyte depletion, followed by one dose of 1928zT2 T cells at
8 5×10^4 cells/kg (Supplementary Figure 11a). After infusion of 1928zT2 T cells, the
9 patient experienced side effects including fever, coughing, skin rashes and
10 hypotension, but not neurological defects. During this trial, the symptoms of the
11 patients did not necessitate ICU stay, vasopressor support or any steroids treatment. A
12 marked elevation of serum IL-6 was detected on day 2 and tocilizumab, an anti-IL-6R
13 monoclonal antibody ⁷, was administered at 4 mg/kg, which resulted in rapid
14 defervescence and complete reversal of symptoms over a period of 4 days; however,
15 the patient's concentration of C-reactive protein (CRP) fluctuated in another pattern,
16 and the highest peak on day 45 was due to urological infection (Supplementary Figure
17 11b). In addition, we monitored the dynamic changes of other cytokines
18 (Supplementary Figure 11c). B-ALL cells comprised approximately 15% of the
19 PBMC and more than 70% of the BM before treatment (Figure 5a), but decreased to
20 an undetectable level on day 22 (Figure 5b). Complete remission (CR) was formally
21 diagnosed on day 51 (Figure 5c). Percentage changes of 1928zT2 T cells and CD19⁺
22 B-ALL cells in PBMC (Figure 5d) and BM (Figure 5e) were depicted. On day 22, the

percentages of $CD4^+CD8^-$, $CD62L^+CD25^-$ and $CD45RO^+CD45RA^-$ populations in 1928zT2 T cells were higher than those in wild-type T cells in both PBMC and BM (Supplementary Figure 12a). Interestingly, the elevation of $CD4^+$ 1928zT2 T cell percentage was concomitant with the increase of the percentage of 1928zT2 T cells in both PBMC and BM (Supplementary Figure 12b and 12c). In addition, the percentages of $CD25^+$ 1928zT2 T cells were negatively correlated with the percentages of 1928zT2 T cells in PBMC (Supplementary Figure 12d), while CD62L expression on 1928zT2 T cells in PBMC remained high until the B-ALL cells were eradicated (Supplementary Figure 12e). 1928zT2 T cells were detected in cerebrospinal fluid, indicating the potential role of 1928zT2 T cells in eradicating leukemia in the central nervous system (Supplementary Figure 12f). The clinical results here indicated that small dosages of 1928zT2 T cells could kill leukemic cells with potent efficacy.

Discussion

Although CD19-specific CAR T cell immunotherapy has achieved promising results in clinical trials, efficacy of other CAR T cells needs improvement, especially those for solid tumors. The development of CARs from the first generation (with only the CD3 ζ signaling domain) to the second generation (with CD3 ζ and a costimulatory domain) has revealed a critical role of costimulatory molecules in CARs. Incorporation of novel costimulatory molecules into CARs should be a feasible strategy to improve their efficacy.

1 Interestingly, the innate immune receptor TLR2 is expressed on activated T cells as
2 a costimulatory receptor¹⁰. Innate immunity serves as an immediate defense
3 mechanism against pathogenic organisms. Adaptive immunity plays a delayed but
4 more delicate role in controlling diseases and forming memories. The two types of
5 immunity are not independent of each other and the interaction between them is
6 critical to the outcome of overall immune response. CARs with combinatorial
7 moieties of both adaptive and innate immunity may exert dual functions of mobilizing
8 innate responses as well as promoting T cell expansion, antigen-specific reaction, T
9 cell contraction and memory formation. We demonstrated that 1928zT2 and m28zT2
10 T cells were more potent at killing CD19⁺ leukemia cells and mesothelin⁺ lung cancer
11 cells in vitro. TLR2 signaling in CD8 T cells can augment T-bet protein levels and
12 IFN- γ , perforin and granzyme B expression through the mTOR pathway, and
13 contribute to the efficient clearance of intracellular pathogens³³, reminiscent of the
14 results we provide here. To decipher other mechanisms of TLR2-enhanced antitumor
15 efficacy, we found the genes upregulated by TLR2 costimulation, most of which were
16 associated with cell adhesion, synaptic transmission and migration. Therefore, TLR2
17 costimulation can serve as an innate immune moiety to enhance the effector functions
18 and promote adhesion and immune synaptic signaling between CAR T cells and target
19 cells.

20 In our pilot study, only 5×10^4 /kg of 1928zT2 T cells expanded to a population that
21 constituted approximately 40% of lymphocytes in patient PBMC, indicating the
22 remarkable expansion capacity of 1928zT2 T cells. Since correlation between infused

1 CAR T cell dose and incidence and severity of cytokine release syndrome (CRS) has
2 been suggested^{26, 37, 38}, it should be beneficial to reduce the CAR T cell dose. In
3 addition, while large numbers of CAR T cells can be generated, a reduction in CAR-T
4 cell dose may reduce the time for producing enough CAR T cells for therapy and
5 manufacturing cost, and provide capacity advantages, which will be important for
6 widespread adoption of CAR-T cell therapies. Nevertheless, other patients in our
7 clinical trial showed less severe side effects and none needed tocilizumab treatment
8 (data unpublished). Therefore, our TLR2 construct should be potent but without
9 increased incidence and severity of CRS. We provided an unprecedentedly detailed
10 dynamic changes of CAR T cells and B-ALL cells in the patient, a pattern which
11 could be reference for other patients to predict CR. However, results of our ongoing
12 clinical trial will be needed to further determine the safety and efficacy of TLR2
13 co-stimulation in larger patient cohorts.

14 CAR T cells for solid tumors face five major challenges — trafficking, tumor
15 recognition and killing, proliferation and persistence, counteract of microenvironment
16 and control of CAR T cells³⁹, and the former four of which can be tackled by
17 optimization of costimulations in CARs. In our A549-GL metastatic xenograft mouse
18 models (Supplementary Figure 9a), the tumors in the m28z group were larger than
19 that not only in the m28zT2 group but also in the GFP group. Conversely, we
20 demonstrated that m28z T cells were able to kill A549-GL cells in vitro (Figure 4a)
21 and suppress the growth of A549-GL in subcutaneous xenograft models (Figure 4c).
22 The differences of the results from these experiments show the importance of

validating the efficacies of cellular immunotherapies in multiple models. The size of tumors is determined by the combination effects of tumor growth and tumor killing in vivo. It is possible that inflammatory factors secreted by m28z T cells or tumor cells, and cell debris during killing stimulated and recruited mouse tumor-supportive cells, such as MDSCs^{40, 41} that easily access to and accumulate in uterus^{42, 43}. Then the tumor-supportive cells promoted the expansion of A549-GL cells and overcame the anti-tumor effects of m28z T cells⁴⁴. However, m28zT2 T cells eliminated A549-GL cells efficiently enough so that tumors hardly formed and grew in NSI mice. Of course, further studies were required to test this hypothesis in the future.

To summarize, our data demonstrate that TLR2 incorporation can generate more powerful CARs, which have superior anti-tumor efficacies in vitro and in vivo and can be used in patients. These findings reveal an important new strategy for CAR design which bridges the conventional costimulation of adaptive immunity with unconventional innate immune signaling.

Acknowledgements

We sincerely thank Dr. Robert Weinkove at Malaghan Institute of Medical Research for his help in revising the manuscript. This study was supported by National Natural Science Foundation of China (81522002), the Strategic Priority Research Program of the Chinese Academy of Sciences (XDB19030205), the Natural Science Fund for Distinguished Young Scholars of Guangdong Province (2014A030306028), the Guangdong Provincial Applied Science and Technology Research& Development

Program (2016B020237006), the Guangdong Provincial Outstanding Young Scholars Award (2014TQ01R068), the Frontier and key technology innovation special grant from the Department of Science and Technology of Guangdong province, (2015B020227003, 2014B020225005, 2016B030229006), the Guangdong Provincial Research and Commercialization Program (2014B090901044), and the Guangzhou Science Technology and Innovation Commission Project (201504010016). A patent application has been filed for the TLR2 incorporated CAR vector, and P. L. and Y. L. are listed as authors on this application. The clinical trial is being sponsored by Guangdong Zhaotai Invivo Biomedicine Co. Ltd. and Hunan Zhaotai Yongren Medical Innovation Ltd.

Conflict of Interest Statement

The authors declare no competing financial interests.

Figure legends

Figure 1. 1928zT2 T cells showed enhanced effector functions compared with 1928z T cells in vitro. **(a)** Killing percentages of K562-GL, K562-CD19-GL, NALM6-GL and REH-GL cells by GFP T, 1928z T and 1928zT2 T cells after 18h coculture in vitro. E: T ratios mean the ratios of absolute number of CAR T cells to target cells. The GFP percentages of CAR T cells were equalized by non-transduced T cells from the same donor. Results are representative of at least three independent experiments with T cells from different healthy donors. **(b)** IL-2, IFN- γ , GM-CSF and TNF α

1 concentration in the supernatant of 24h coculture of the indicated T cells with
2 K562-GL or K562-CD19-GL cells (E: T=1, 2×10^5 cells each). (c) Enhanced
3 expansion and serial killing potency of 1928zT2 T cells compared with 1928z T cells.
4 1928z T and 1928zT2 T cells were serially co-cultured with NALM6 cells at 1: 4 E/T
5 ratio, the numbers of 1928z T cells and 1928zT2 T cells were counted (left), and
6 percentages of CD19⁺ NALM6 cells (right) were detected by flow cytometry, every
7 24h for three times. Error bars denote s.e.m and results were compared through
8 two-sided unpaired *t*-test. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

9 **Figure 2.** 1928zT2 T cells showed superior anti-leukemia efficacy in vivo. (a)
10 Bioluminescence imaging of NALM6-GL-burdened mice treated with GFP T, 1928z
11 T or 1928zT2 T cells. 1×10^6 NALM6-GL cells were injected intravenously into NSI
12 mice, and on day 2 and 7, 5×10^6 GFP T, 1928z T or 1928zT2 T cells were injected
13 intravenously into each NALM6-GL-NSI mice (n=5). On day 8, 14 and 19,
14 bioluminescence imaging was conducted. (b) The percentages of CD19⁺ (NALM6-GL)
15 cells in the PBMC of mice on day 14. (c) Survival analysis of mice treated with GFP
16 T, 1928z T or 1928zT2 T cells (n=10). 2×10^5 NALM6 cells were intravenously
17 injected into NSI mice, and two days later, 2×10^6 T cells were injected (GFP%
18 normalized to 10%), and survival of mice was observed. Error bars denote s.e.m and
19 groups were compared through two-sided unpaired *t*-test. * $P < 0.05$, ** $P < 0.01$, ***
20 $P < 0.001$.

21 **Figure 3.** TLR2 costimulation up-regulated genes associated with cell adhesion,
22 synaptic transmission, infiltration and mTOR pathway. (a) Heatmap of RNA

sequencing results showing genes specifically up-regulated in 1928zT2 T cells after stimulation (+sti) (left). These genes were associated with cell adhesion, synapse components and transmission and cell invasion of T cells (right). (b) Relative expression of genes associated with cell adhesion (*CD24*, *CDH2*, *TYRO3*, *CXADR*), T cell infiltration (*MMP2*), synaptic transmission (*CADPS2*), TLRs pathway (*IRF3* and *IRF7*) and mTOR pathway (*T-bet*, *mTOR*, *4E-BP1* and *p70S6K*). (c) Transwell cell migration assay indicated an improved capacity of 1928zT2 T cells to degrade and transmigrate Matrigel. T cell culture medium was used as a chemoattractant in the lower chamber. T cells were cultured in the insert with or without coated Matrigel for 24 h, and the cells that transmigrated to the lower chamber were counted. The percentage of invasion was calculated as follows: (mean of cells migrating through the Matrigel chamber membrane/mean of cells migrating through the control insert membrane) $\times 100$. Error bars denote s.e.m and groups were compared through two-sided unpaired *t*-test. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

Figure 4. m28zT2 T cells showed enhanced antitumor efficacy compared with m28z T cells. (a) Killing percentages of mesothelin⁺ A549-GL and mesothelin⁻ H23-GL cells after 18h coculture with GFP T, m28z T or m28zT2 T cells at the indicated ratios. (b) IL-2, IFN- γ , GM-CSF and TNF- α concentration in the supernatant of 24h coculture of GFP T, m28z T or m28zT2 T cells with A549-GL cells. E: T=1:1, 2×10^5 cells each. (c) Weight of subcutaneous A549-GL tumors from NSI mice treated with no T cells, GFP T, m28z T or m28zT2 T cells. On day 0, 2×10^5 A549-GL cells were injected subcutaneously into NSI mice, and on day 7 and 9, no T, 5×10^6 GFP T, m28z

1 T or m28zT2 T cells were injected through the tail vein into each mouse (n=6). On
 2 day 43, mice were sacrificed and tumors were analyzed. Error bars denote s.e.m and
 3 groups were compared through two-sided unpaired *t*-test. * $P < 0.05$, ** $P < 0.01$, ***
 4 $P < 0.001$.

5 **Figure 5.** 1928zT2 T cells showed potent efficacy in a patient with relapsed B-ALL.
 6 (a) FACS analysis of B-ALL cells in PBMC and BM from the patient on the day
 7 before infusion of 1928zT2 T cells. (b) B-ALL cells were eradicated from PBMC and
 8 BM of the patient on day 19 and 22 respectively. (c) Wright-Giemsa staining of BM
 9 biopsies of the patient on day 18 before infusion (left) and on day 51 post-infusion
 10 (right) of 1928zT2 T cells. Scale bar = 10 μ m. (d) Dynamic changes of the percentages
 11 of CD19⁺ cells and 1928zT2 T cells in PBMC of the patient. (e) Dynamic changes of
 12 the percentages of CD19⁺ cells and 1928zT2 T cells in BM of the patient.

13
 14

15 References

- 16 1. Hombach A, Abken H. Costimulation tunes tumor-specific activation of redirected T cells in
 17 adoptive immunotherapy. *Cancer Immunol Immunother* 2007 May; **56**(5): 731-737.
- 18 2. Kowolik CM, Topp MS, Gonzalez S, Pfeiffer T, Olivares S, Gonzalez N, *et al.* CD28 costimulation
 19 provided through a CD19-specific chimeric antigen receptor enhances in vivo persistence and
 20 antitumor efficacy of adoptively transferred T cells. *Cancer Res* 2006 Nov 15; **66**(22):
 21 10995-11004.
 22
- 23 3. Milone MC, Fish JD, Carpenito C, Carroll RG, Binder GK, Teachey D, *et al.* Chimeric receptors
 24 containing CD137 signal transduction domains mediate enhanced survival of T cells and
 25 increased antileukemic efficacy in vivo. *Mol Ther* 2009 Aug; **17**(8): 1453-1464.
 26
- 27 4. Long AH, Haso WM, Shern JF, Wanhainen KM, Murgai M, Ingaramo M, *et al.* 4-1BB
 28 costimulation ameliorates T cell exhaustion induced by tonic signaling of chimeric antigen
 29

- 1 receptors. *Nat Med* 2015 Jun; **21**(6): 581-590.
- 2
- 3 5. Pule MA, Straathof KC, Dotti G, Heslop HE, Rooney CM, Brenner MK. A chimeric T cell antigen
4 receptor that augments cytokine release and supports clonal expansion of primary human T
5 cells. *Mol Ther* 2005 Nov; **12**(5): 933-941.
- 6
- 7 6. Song DG, Ye Q, Poussin M, Harms GM, Figini M, Powell DJ, Jr. CD27 costimulation augments
8 the survival and antitumor activity of redirected human T cells in vivo. *Blood* 2012 Jan 19;
9 **119**(3): 696-706.
- 10
- 11 7. Shen CJ, Yang YX, Han EQ, Cao N, Wang YF, Wang Y, *et al.* Chimeric antigen receptor
12 containing ICOS signaling domain mediates specific and efficient antitumor effect of T cells
13 against EGFRvIII expressing glioma. *J Hematol Oncol* 2013; **6**: 33.
- 14
- 15 8. Zhang BL, Qin DY, Mo ZM, Li Y, Wei W, Wang YS, *et al.* Hurdles of CAR-T cell-based cancer
16 immunotherapy directed against solid tumors. *Science China Life sciences* 2016 Apr; **59**(4):
17 340-348.
- 18
- 19 9. Brown CE, Alizadeh D, Starr R, Weng L, Wagner JR, Naranjo A, *et al.* Regression of
20 Glioblastoma after Chimeric Antigen Receptor T-Cell Therapy. *N Engl J Med* 2016 Dec 29;
21 **375**(26): 2561-2569.
- 22
- 23 10. Komai-Koma M, Jones L, Ogg GS, Xu DM, Liew FY. TLR2 is expressed on activated T cells as a
24 costimulatory receptor. *P Natl Acad Sci USA* 2004 Mar 2; **101**(9): 3029-3034.
- 25
- 26 11. Cottalorda A, Mercier BC, Mbitikon-Kobo FM, Arpin C, Teoh DY, McMichael A, *et al.* TLR2
27 engagement on memory CD8(+) T cells improves their cytokine-mediated proliferation and
28 IFN-gamma secretion in the absence of Ag. *Eur J Immunol* 2009 Oct; **39**(10): 2673-2681.
- 29
- 30 12. Sinnott BD, Park B, Boer MC, Lewinsohn DA, Lancioni CL. Direct TLR-2 Costimulation Unmasks
31 the Proinflammatory Potential of Neonatal CD4+ T Cells. *J Immunol* 2016 Jul 1; **197**(1): 68-77.
- 32
- 33 13. Salerno F, Guislain A, Cansever D, Wolkers MC. TLR-Mediated Innate Production of
34 IFN-gamma by CD8+ T Cells Is Independent of Glycolysis. *J Immunol* 2016 May 1; **196**(9):
35 3695-3705.
- 36
- 37 14. Cottalorda A, Verschelde C, Marçais A, Tomkowiak M, Musette P, Uematsu S, *et al.* TLR2
38 engagement on CD8 T cells lowers the threshold for optimal antigen-induced T cell activation.
39 *Eur J Immunol* 2006 Jul; **36**(7): 1684-1693.
- 40
- 41 15. Mercier BC, Cottalorda A, Coupet CA, Marvel J, Bonnefoy-Berard N. TLR2 engagement on CD8
42 T cells enables generation of functional memory cells in response to a suboptimal TCR signal.
43 *J Immunol* 2009 Feb 15; **182**(4): 1860-1867.
- 44

- 1 16. Quigley M, Martinez J, Huang X, Yang Y. A critical role for direct TLR2-MyD88 signaling in CD8
2 T-cell clonal expansion and memory formation following vaccinia viral infection. *Blood* 2009
3 Mar 5; **113**(10): 2256-2264.
4
- 5 17. Imanishi T, Hara H, Suzuki S, Suzuki N, Akira S, Saito T. Cutting Edge: TLR2 Directly Triggers Th1
6 Effector Functions. *The Journal of Immunology* 2007; **178**(11): 6715-6719.
7
- 8 18. Oberg HH, Ly TT, Ussat S, Meyer T, Kabelitz D, Wesch D. Differential but direct abolishment of
9 human regulatory T cell suppressive capacity by various TLR2 ligands. *J Immunol* 2010 May 1;
10 **184**(9): 4733-4740.
11
- 12 19. Babu S, Blauvelt CP, Kumaraswami V, Nutman TB. Cutting Edge: Diminished T Cell TLR
13 Expression and Function Modulates the Immune Response in Human Filarial Infection. *The*
14 *Journal of Immunology* 2006; **176**(7): 3885-3889.
15
- 16 20. Zhang Y, Luo F, Cai Y, Liu N, Wang L, Xu D, *et al.* TLR1/TLR2 agonist induces tumor regression
17 by reciprocal modulation of effector and regulatory T cells. *J Immunol* 2011 Feb 15; **186**(4):
18 1963-1969.
19
- 20 21. Zhang Y, Luo F, Li A, Qian J, Yao Z, Feng X, *et al.* Systemic injection of TLR1/2 agonist improves
21 adoptive antigen-specific T cell therapy in glioma-bearing mice. *Clin Immunol* 2014 Sep;
22 **154**(1): 26-36.
23
- 24 22. Sandor F, Latz E, Re F, Mandell L, Repik G, Golenbock DT, *et al.* Importance of extra- and
25 intracellular domains of TLR1 and TLR2 in NFkappa B signaling. *J Cell Biol* 2003 Sep 15; **162**(6):
26 1099-1110.
27
- 28 23. O'Neill LA, Bowie AG. The family of five: TIR-domain-containing adaptors in Toll-like receptor
29 signalling. *Nat Rev Immunol* 2007 May; **7**(5): 353-364.
30
- 31 24. Brown V, Brown RA, Ozinsky A, Hesselberth JR, Fields S. Binding specificity of Toll-like
32 receptor cytoplasmic domains. *Eur J Immunol* 2006 Mar; **36**(3): 742-753.
33
- 34 25. Brentjens RJ, Davila ML, Riviere I, Park J, Wang X, Cowell LG, *et al.* CD19-targeted T cells
35 rapidly induce molecular remissions in adults with chemotherapy-refractory acute
36 lymphoblastic leukemia. *Sci Transl Med* 2013 Mar 20; **5**(177): 177ra138.
37
- 38 26. Davila ML, Riviere I, Wang X, Bartido S, Park J, Curran K, *et al.* Efficacy and toxicity
39 management of 19-28z CAR T cell therapy in B cell acute lymphoblastic leukemia. *Sci Transl*
40 *Med* 2014 Feb 19; **6**(224): 224ra225.
41
- 42 27. Grupp SA, Kalos M, Barrett D, Aplenc R, Porter DL, Rheingold SR, *et al.* Chimeric antigen
43 receptor-modified T cells for acute lymphoid leukemia. *N Engl J Med* 2013 Apr 18; **368**(16):
44 1509-1518.

28. Kochenderfer JN, Dudley ME, Carpenter RO, Kassim SH, Rose JJ, Telford WG, *et al.* Donor-derived CD19-targeted T cells cause regression of malignancy persisting after allogeneic hematopoietic stem cell transplantation. *Blood* 2013 Dec 12; **122**(25): 4129-4139.
29. Maude SL, Frey N, Shaw PA, Aplenc R, Barrett DM, Bunin NJ, *et al.* Chimeric antigen receptor T cells for sustained remissions in leukemia. *N Engl J Med* 2014 Oct 16; **371**(16): 1507-1517.
30. Porter DL, Levine BL, Kalos M, Bagg A, June CH. Chimeric antigen receptor-modified T cells in chronic lymphoid leukemia. *N Engl J Med* 2011 Aug 25; **365**(8): 725-733.
31. Dobrovolskaia MA, Medvedev AE, Thomas KE, Cuesta N, Toshchakov V, Ren T, *et al.* Induction of in vitro reprogramming by Toll-like receptor (TLR)2 and TLR4 agonists in murine macrophages: effects of TLR "homotolerance" versus "heterotolerance" on NF-kappa B signaling pathway components. *J Immunol* 2003 Jan 1; **170**(1): 508-519.
32. Ye W, Jiang Z, Li GX, Xiao Y, Lin S, Lai Y, *et al.* Quantitative evaluation of the immunodeficiency of a mouse strain by tumor engraftments. *Journal of hematology & oncology* 2015; **8**: 59.
33. Geng D, Zheng L, Srivastava R, Asprodites N, Velasco-Gonzalez C, Davila E. When Toll-like receptor and T-cell receptor signals collide: a mechanism for enhanced CD8 T-cell effector function. *Blood* 2010 Nov 04; **116**(18): 3494-3504.
34. Boussif O, Lezoualc'h F, Zanta MA, Mergny MD, Scherman D, Demeneix B, *et al.* A versatile vector for gene and oligonucleotide transfer into cells in culture and in vivo: polyethylenimine. *Proc Natl Acad Sci U S A* 1995 Aug 1; **92**(16): 7297-7301.
35. Ye W, Jiang Z, Li GX, Xiao Y, Lin S, Lai Y, *et al.* Quantitative evaluation of the immunodeficiency of a mouse strain by tumor engraftments. *J Hematol Oncol* 2015; **8**: 59.
36. Ho M, Bera TK, Willingham MC, Onda M, Hassan R, FitzGerald D, *et al.* Mesothelin expression in human lung cancer. *Clin Cancer Res* 2007 Mar 1; **13**(5): 1571-1575.
37. Lee DW, Kochenderfer JN, Stetler-Stevenson M, Cui YK, Delbrook C, Feldman SA, *et al.* T cells expressing CD19 chimeric antigen receptors for acute lymphoblastic leukaemia in children and young adults: a phase 1 dose-escalation trial. *Lancet* 2015 Feb 07; **385**(9967): 517-528.
38. Park JH, Geyer MB, Brentjens RJ. CD19-targeted CAR T-cell therapeutics for hematologic malignancies: interpreting clinical outcomes to date. *Blood* 2016 Jun 30; **127**(26): 3312-3320.
39. Lim WA, June CH. The Principles of Engineering Immune Cells to Treat Cancer. *Cell* 2017 Feb 09; **168**(4): 724-740.
40. Sevko A, Umansky V. Myeloid-derived suppressor cells interact with tumors in terms of

- 1 myelopoiesis, tumorigenesis and immunosuppression: thick as thieves. *J Cancer* 2013; **4**(1):
2 3-11.
- 3
- 4 41. Ortiz ML, Lu L, Ramachandran I, Gabrilovich DI. Myeloid-derived suppressor cells in the
5 development of lung cancer. *Cancer Immunol Res* 2014 Jan; **2**(1): 50-58.
- 6
- 7 42. Ostrand-Rosenberg S, Sinha P, Figley C, Long R, Park D, Carter D, *et al.* Frontline Science:
8 Myeloid-derived suppressor cells (MDSCs) facilitate maternal-fetal tolerance in mice. *J Leukoc*
9 *Biol* 2017 May; **101**(5): 1091-1101.
- 10
- 11 43. Zhao H, Kalish F, Schulz S, Yang Y, Wong RJ, Stevenson DK. Unique roles of infiltrating myeloid
12 cells in the murine uterus during early to midpregnancy. *J Immunol* 2015 Apr 15; **194**(8):
13 3713-3722.
- 14
- 15 44. Umansky V, Blattner C, Gebhardt C, Utikal J. The Role of Myeloid-Derived Suppressor Cells
16 (MDSC) in Cancer Progression. *Vaccines* 2016 Nov 03; **4**(4).
- 17
- 18